

# NANOTRANSPORT -

## *Transport of Engineered Nanoparticle (ENP) Aerosols in the Workplace Environment and its <sup>L</sup> Consequences*

### Recommendations to European Commission regarding

- *Test aerosols for nano-toxicology studies*
- *Testing of filters and protective equipment in the workplace*
- *Metrology of nano-aerosols*
- *Open questions: Research priorities*



# Essential constraints of the study

- *Study limited to airborne NP („nano-aerosols“)*
- *Study concentrated only on the aspect of **NP size** as a key metric,*
- *the **evolution of „size“ with time** spent in the aerosol phase („history“) **between source and receptor** (i.e. target organism), and its implications for measurement, testing, abatement, hazard evaluation.*

## Key conclusions from the study

- *Nano-aerosols evolve considerably with time: their average size increases while concentration decreases*
- ***Natural background aerosols** are effective scavengers for NP*
- *Thus, NP will (usually) be physically/chemically present in size classes other than the one in which they were originally emitted.*
- ***Time scale** for size evolution depends on concentration - on the order of a few min.*
- *Filtration efficiency for „true“ NP not an issue - focus on filter performance at MPPS.*



# Test aerosols for *nano-toxicology studies*

- „Aged“ NP aerosols and primary sources of (unagglomerated) NP equally relevant for risk assessment in workplace environments
- **Scenario I:** direct exposure to primary NP aerosol
  - only for people standing **very close to source**
- **Scenario II:** indirect exposure via inhalation of aged/attached NP
  - Includes secondary NP aerosols due to re-suspension from surfaces, which are always agglomerated



# Test aerosols for *nano-toxicology studies* (2)

## **Characteristics of „aged“ nano-aerosols:**

- Typical size range for auto-agglomeration: (20) - 200 nm
- Typical size range for attachment to background aerosol:  $0.1 < d < 1 \mu\text{m}$   
(number concentration in the coarse aerosol mode usually too low)
- „Aging“ in the aerosol phase by Brownian collisions usually leads to **relatively weak agglomerate structures**
- **Consequences:**
  - Strongly recommend **generating test particles via aerosol phase** (incl. aging, scavenging), rather than instillation of powders or liquid dispersions
  - **Control of particle structure** (as well as surface chemistry) critical for toxicological studies (not only size), because influence of agglomerate strength, structure, etc. on toxicity not well understood
  - Control of structure, “history” of particles much better done **via aerosol processes** involve aerosol technologists in these studies!



# Testing of filters & **protective equipment** in the workplace

- *Filtration efficiency for “true“ NP (2-10 nm) generally sufficient*
- *Need to consider effective protection against scavenged NP*
- *Most critical size range around filter MPPS: 80 - 200 nm*
- *Effective testing equipment and aerosol generation in this range most important (SMPS+CPC)*
- *Physical/chemical composition of test aerosol not relevant for filtration, only equivalent size*



# ***Metrology of nano-aerosols in workplace***

- Aged NP aerosols *not recognizable by a specific particle size range* (e.g. "<20 nm")
- Not sufficient to look for NP in the "nano size range" where they may have originally been generated/emitted
- Not sufficient to assess workplace situation via simple size distribution measurements - additional ENP concentrations are often marginal
- ... unless an analytical technique is *highly selective*
- *Alternative: aerosol dynamic models to predict ENP hazard in spec. scenarios*
  - characterize *emission sources* (strength, primary size range)
  - *identify release mechanisms* for each specific ENP
- Need capability for *species-specific tracing* of scavenged/agglom. NP
- Need to validate (existing!) aerosol dynamic models for workplace scenarios
- Need technique for measuring agglomerate strength



# Open questions & Research priorities

- Characterize **release mechanisms and sources** of hazardous ENP types (source strength, size range, particle structure) for specific scenarios
- Develop **aerosol generators** for relevant classes of release mechanisms and species for use in toxicological studies
  - incl. controlled/variable NP structure, surface chemistry
- Validate & refine **aerosol dynamic models** for predicting ENP evolution between source & receptor for specific scenarios
- Develop **species-specific real-time detection methods**
  - real-time chemical detection of attached NP aerosols
  - agglomerate strength
  - Surface composition
  - ...
- *Involve more aerosol technologists and their tools in toxicology studies!*

